

“BLUE GEL” ANESTHETIC

For Use by Licensed Professionals Only.

This product contains 4% lidocaine and 2% tetracaine, together with a vascular constrictor which reduces swelling, bruising and bleeding. This makes it the anesthetic of choice for fast onset and reduction of areas, such as lips or inner arms, where bruising and bleeding are a problem.

FOR EXTERNAL USE ONLY: Temporarily relieves pain and swelling associated with tattooing, permanent makeup, piercing or other pain sensitive procedures. Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool, dark place, *out of reach of children*. Discard by expiration date.

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline and seek medical attention. **DO NOT SWALLOW.** If you should accidentally swallow this product seek medical attention immediately. Do not use if pregnant or nursing.

DURING PROCEDURE ANESTHETIC

2-5 minutes

Use this product during painful procedures to continue the anesthetic effects of your pre-procedure topical anesthetics, or to deaden broken skin. *It will have no effect if applied to unbroken skin.*

Always wear gloves while applying. Apply to the procedure area with an applicator, cover with plastic and wait 2-5 minutes or until anesthetic effect occurs.

Rinse off thoroughly before continuing your procedure so that the glycerin base does not saturate the skin. **DO NOT REAPPLY** this product to the same areas because it can cause the skin to become so numb, you could harm the tissue and the tattoo. If you need more anesthetic effect before you have finished your procedure, use *SJ3* or *Vasocaine* to continue the anesthetic effect.

Blue Gel will keep the skin anesthetized and constricted for up to two hours.

SUGGESTED APPLICATION

DERMAL SOURCE
1017 SW Morrison Street, #210
Portland, Oregon 97205
Phone: 866-568-3223, Fax: 503-973-5433
www.dermalsouce.com



05/10