

MAXIMUM ZONE 2

For Use by Licensed Professionals Only

This product contains 5% Lidocaine and a vascular constrictor, which reduces swelling, bruising, and bleeding. This makes it the choice for fast onset and reduction of areas, such as lips or inner arms, where bruising and bleeding are a problem.

FOR EXTERNAL USE ONLY: Temporarily relieves pain, itching or swelling associated with anorectal disorders or other pain sensitive procedures such as laser, electrolysis, tattooing or piercing. Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool, dark place, *out of reach of children*.

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline and seek medical attention. **DO NOT SWALLOW.** If you should accidentally swallow this product, seek medical attention immediately. Do not use if pregnant or nursing.

DURING PROCEDURE ANESTHETIC

2-5 minutes

Use this product during painful procedures to continue the anesthetic effects of your pre-procedure topical anesthetics, or to anesthetize broken skin. *It will have no effect if applied to intact skin.*

Always wear gloves while applying this product. Apply to the procedure area with an applicator, occlude with plastic and wait 2-5 minutes or until anesthetic effect occurs.

Rinse off thoroughly before continuing your procedure so that the glycerin base does not saturate the skin. **DO NOT REAPPLY** this product to the same areas because it can cause the skin to become so anesthetized that you could harm the tissue. If you need more anesthetic effect before you have finished your procedure, use **Superior Super Juice** or **Five-Star Vasocaine** to continue the anesthetic effect.

Maximum Zone 2 will keep the skin anesthetized and constricted for up to two hours.

SUGGESTED APPLICATION

DERMAL SOURCE

2319 NE Glisan Street, Portland, Oregon 97232, U.S.A.

Phone: +1 866-568-3223, Fax: 503-595-0293

www.dermalsource.com



MAXIMUM ZONE 2

For Use by Licensed Professionals Only

This product contains 5% Lidocaine and a vascular constrictor, which reduces swelling, bruising, and bleeding. This makes it the choice for fast onset and reduction of areas, such as lips or inner arms, where bruising and bleeding are a problem.

FOR EXTERNAL USE ONLY: Temporarily relieves pain, itching or swelling associated with anorectal disorders or other pain sensitive procedures such as laser, electrolysis, tattooing or piercing. Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool, dark place, *out of reach of children*.

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline and seek medical attention. **DO NOT SWALLOW.** If you should accidentally swallow this product, seek medical attention immediately. Do not use if pregnant or nursing.

DURING PROCEDURE ANESTHETIC

2-5 minutes

Use this product during painful procedures to continue the anesthetic effects of your pre-procedure topical anesthetics, or to anesthetize broken skin. *It will have no effect if applied to intact skin.*

Always wear gloves while applying this product. Apply to the procedure area with an applicator, occlude with plastic and wait 2-5 minutes or until anesthetic effect occurs.

Rinse off thoroughly before continuing your procedure so that the glycerin base does not saturate the skin. **DO NOT REAPPLY** this product to the same areas because it can cause the skin to become so anesthetized that you could harm the tissue. If you need more anesthetic effect before you have finished your procedure, use **Superior Super Juice** or **Five-Star Vasocaine** to continue the anesthetic effect.

Maximum Zone 2 will keep the skin anesthetized and constricted for up to two hours.

SUGGESTED APPLICATION

DERMAL SOURCE

2319 NE Glisan Street, Portland, Oregon 97232, U.S.A.

Phone: +1 866-568-3223, Fax: 503-595-0293

www.dermalsource.com

