

“PRO PLUS” TOPICAL ANESTHETIC

For Use by Licensed Professionals Only.

This pre-deadener was developed by a pharmacist familiar with our industries. Previous versions of this anesthetic have been used successfully for years. This recent alteration has improved its speed and depth of anesthesia. This cream contains 2% tetracaine and 4% lidocaine, has a pH of 7 and is *safe for eyeliner*.

FOR EXTERNAL USE ONLY ON INTACT SKIN: Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool dark place, *out of the reach of children*.

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline and contact an eye care physician. **DO NOT SWALLOW.** If accidentally swallowed, seek medical help immediately.

SUGGESTED APPLICATION TIMES

EYEBROWS

15- 20 minutes

For best results, lightly exfoliate with *Skin Resurfacer* prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing.

EYELINER

15- 20 minutes

Fill the eyes with liquid eye gel to coat the cornea. With the client's eyes closed, apply a thin layer of cream with a small applicator across the lash line. Remove anesthetic before beginning procedure. Following the first pass of tattooing, sparingly apply Zone 2 or Blue Gel to control swelling and maintain anesthesia for up to two hours.

LIPS

15-20 minutes

For best results, lightly exfoliate with *Skin Resurfacer* prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing. While some relief may be provided with topical anesthetics, for total anesthesia, you must provide a nerve block performed by a physician. *Do not use inside of the mouth.* Use *SJ3* during procedure to continue anesthesia, or either *Blue Gel*, *Vasocaine* or *Zone 2* to continue anesthesia and constrict blood vessels.

BODY SKIN

90-120 minutes

Prior to tattooing or piercing: Lightly exfoliate with *Skin Resurfacer*. Gently massage into skin and then apply a medium coating over the top. In order to prevent absorption into clothing and retain effective anesthesia, cover with an occlusive dressing. Thicker skin requires longer time to absorb the cream than thinner skin. Use *SJ3* during procedure to continue anesthesia, or either *Blue Gel*, *Vasocaine* or *Zone 2* to continue anesthesia and constrict blood vessels.

PIERCING

15- 20 minutes

Apply to eyebrows, navels and nipples. Avoid genital areas. Thicker skin requires longer pre-treating time. Lightly exfoliate with *Skin Resurfacer*. Gently massage into skin, then apply a medium coating on top. Cover with occlusive dressing. Wipe away immediately prior to piercing. Use *Blue Gel* or *Zone 2* during jewelry insertion to continue anesthesia and reduce swelling.

DERMAL SOURCE
1017 SW Morrison Street, #210
Portland, Oregon 97205
Phone: 866-568-3223, Fax: 503-973-5433
www.dermalsouce.com



05/10